



TUSSOCK – green skink – ti kouka, young plains ecosystem

Food for native birds:

F = Fruit

S = Bird Seed

N = Nectar

B = Bud/foliage

I = Insects

For lizards: L = fruit

Plant Tolerances

■ = tolerates or needs

□ = intolerant

1/2 = tolerant of some

* = to establish, protect from frost
t = toxic for toddlers

Staging

1 = 1st structural

2 = 2nd year

3 = only after canopy closure

PLANT LISTS

Selected from vegetation natural to these droughty & shallow Selwyn soils

SMALL TREES (> 3 m)

<i>Cordyline australis</i>	ti kouka, cabbage tree	F,N,I
<i>Kunzea ericoides</i>	kanuka	I
<i>Sophora microphylla</i>	South Island kowhai	N,I

SHRUBS & SCRAMBLERS (< 3 m)

<i>Carmichaelia australis</i>	makaka, NZ broom	S,I
<i>Clematis afoliata</i>	leafless clematis	S,I
<i>Clematis quadibracteolata/marāta</i>	clematu	I
<i>Coprosma acerosa</i>	sand coprosma	L
<i>Coprosma propinqua</i>	mikimiki, mingimingi	F,L
<i>Coriaria sarmentosa</i>	shrub tutu	F,I
<i>Discaria toumatou</i>	matagouri	I
<i>Melicytus alpinus</i>	porcupine shrub	F,L,I
<i>Muehlenbeckia astonii</i>	shrub pohuehue	F,L,I
<i>Muehlenbeckia complexa</i>	scrambling pohuehue	F,L,I
<i>Olearia adenocarpa</i>	Canterbury shrub daisy	S,I
<i>Ozothamnus leptophyllus (Cassinia)</i>	tauhinu, cottonwood	S,I
<i>Parsonsia heterophylla</i>	kaiwhiria, New Zealand jasmine	I
<i>Pomaderris ericifolia</i>	tauhinu	S,I
<i>Rubus squarrosus</i>	leafless lawyer	F,L,I
<i>Sophora prostrata</i>	prostrate kowhai	N,I

Tolerances

Food	sun shade	wet dry	wind	Stages
F,N,I	■ 1/2	■ ■	■	1
I	■ □	□ ■	■	1
N,I	■ 1/2	■ ■	t	1

Underlays: Greywacke river stones (2-100mm rounded) & sand on more stones.

